

ERRATUM Open Access

CrossMark

Erratum: Less functional variants of TLR-1/-6/-10 genes are associated with age

Lutz Hamann^{1*}, Juozas Kupcinskas^{2,3}, Luis C. Almanza Berrocal¹, Jurgita Skieceviciene³, Andre Franke⁴, Ute Nöthlings^{5,6} and Ralf R. Schumann¹

After publication of this study [1], the authors noticed that the spelling of author Luis C Berrocal Almanza's name was incorrect. The original version of this article has been updated to correct this. The publisher apologises for any inconvenience caused.

Author details

¹Institute of Microbiology and Hygiene, Charité University Medical Center Berlin, Charitéplatz 1, Berlin 10117, Germany. ²Department of Gastroenterology, Lithuanian University of Health Sciences, Kaunas, Lithuania. ³Institute for Digestive Research, Lithuanian University of Health Sciences, Kaunas, Lithuania. ⁴University Hospital Schleswig Holstein, Campus Kiel, Schittenhelmstr. 12, Kiel, Germany. ⁵Popgen Biobank, Institute for Experimental Medicine, Christian-Albrechts-University Kiel, Niemansweg 11, Kiel, Germany. ⁶Present address: Department of Nutrition and Food Sciences, Rheinische Friedrich-Wilhelms-University Bonn, Endenicher Allee 11-13, Bonn 53115, Germany.

Received: 4 August 2015 Accepted: 4 August 2015 Published online: 11 August 2015

Reference

 Hamann L, Kupcinskas J, Berrocal Almanza LC, Skieceviciene J, Franke A, Nöthlings U, et al. Less functional variants of TLR-1/-6/-10 genes are associated with age. Immun Ageing. 2015;12:7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit





¹Institute of Microbiology and Hygiene, Charité University Medical Center Berlin, Charitéplatz 1, Berlin 10117, Germany

